

Breast Non Feeding: Main Cause of Neonatal Hyperbilirubinemia in Areas Adjoining Shri Ram Murti Smarak Institute of Medical Sciences, A Tertiary Care Teaching Hospital, Bareilly

Shaheena Kamal¹, Kauser Sayedda², Quazi Shahir Ahmed²

¹ Department of Biochemistry

² Department of Pharmacology

Shri Ram Murti Smarak Institute of Medical Sciences (SRMSIMS), Bareilly(UP), India

Correspondence to:

Dr. Shaheena Kamal
(shaheena_ara@yahoo.com)

Received: 02.02.2012

Accepted: 13.06.2012

DOI: 10.5455/njppp.2012.2.108-112

ABSTRACT

Aims & Objective: The purpose of this study was to determine the occurrence, etiological & other associated factors of neonatal hyperbilirubinemia in Shri Ram Murti Smarak Institute of Medical Sciences, Bareilly. Jaundice is a common problem in neonatology. Early recognition of the cause of jaundice is very important as delay in management may lead to serious complications or even death.

Materials and methods: In present study, newborns with jaundice were evaluated during a six months period between July – December 2011. 124 newborns with jaundice were enrolled in the study. Data regarding demographic profile of new born, physical examination and laboratory investigations gathered and analysed to interpretate the common etiologies giving rise to neonatal hyperbilirubinemia.

Results: Out of 124 cases of neonatal hyperbilirubinemia, 24 cases were of physiological jaundice, breast non feeding jaundice, breast milk jaundice, jaundice due to prematurity and pathological jaundice comprised the rest 100 as 84, 5, 5 & 6 cases respectively. Pathological causes for jaundice included neonatal sepsis (2cases), neonatal hypothyroidism (2cases), congenital biliary atresia (1case) and ABO incompatibility (1case).

Conclusion: Present study concludes that breast nonfeeding or inadequate feeding jaundice forms the bulk of cases of neonatal hyperbilirubinemia in this region, followed by breast milk jaundice, prematurity jaundice and pathological causes.

KEY WORDS: Neonatal Hyperbilirubinemia; Breast Non Feeding; Physiological

INTRODUCTION

Hyperbilirubinemia is a common problem during the neonatal period occurring in upto 60% of term and 80% of preterm babies in the first week of life.^[1,2] Some of the most common causes of neonatal jaundice include physiological jaundice, breast feeding or non feeding jaundice, breast milk jaundice, prematurity leading to jaundice & various pathological causes like haemolytic disease, liver dysfunction, neonatal sepsis, deficiency of G6PD enzyme, hypothyroidism and rare conditions such as gilbert's syndrome etc.^[3,4] Extreme hyperbilirubinemia is rare, however, if left untreated especially in premature infant, indirect hyperbilirubinemia may lead to kernicterus, a serious neurological problem and social & economic burden on the patient's family & society.^[5,6] Elevation of direct bilirubin constitute the pathological causes of jaundice & should be promptly treated either by medical or surgical means.^[7,8]

Neonatal jaundice is related to breast feeding in two primary clinical situations: firstly, newborns who receive inadequate breast feeding & have high concentration of indirect bilirubin during the first post natal week (breast non feeding jaundice).^[9] Rise in indirect bilirubin in this type is due to enhancement of enterohepatic circulation. Moreover, presence of large amounts of bilirubin in meconium & delay in emptying of meconium have been shown to contribute to an increase in serum bilirubin levels in early days of life further increasing intestinal biliary absorption.^[10,11] The jaundice in this case can usually be ameliorated by frequent breast feeding sessions of sufficient duration to stimulate adequate milk production.^[12] Passage of baby through vagina during birth helps stimulate milk production in the mother's body, so infants born by caesarean section are at higher risk for this condition. Secondly, breast fed infant who experience prolonged unconjugated hyperbilirubinemia known as breast milk jaundice.^[9] Several hypotheses have been proposed for breast milk jaundice, including the presence of UDP glucuronosyltransferase inhibitor, β glucuronidase or yet unidentified

factor in human milk that could inhibit excretion of bilirubin & results in hyperbilirubinemia. There is increase in enterohepatic circulation of bilirubin in this type also and may be attributed to increased levels of epidermal growth factor (EGF) in breast milk.^[13]

Consideration of all these etiologies is essential in evaluating neonates with jaundice. Complete history of newborn, family history, sibling history, complete physical examination and laboratory investigations are key points in managing these patients.^[14]

The aim of our study is to determine the underlying aetiologies of neonatal hyperbilirubinemia and to explore most common cause of neonatal jaundice in areas adjoining this tertiary care teaching hospital.

MATERIALS AND METHODS

We limited our study in 124 neonates. Out of these, 82 were inborn patients & 42 were outborn. Patients' characteristics & general data were documented including age, birth weight, age at onset, type of delivery, type of feeding, history of jaundice in sibling. All cases were thoroughly examined on admission. Various investigations were carried out including serum bilirubin total & conjugated by Jendrasik and Graf method, liver function tests (aspartate aminotransferase, alkaline phosphatases) by IFCC method by fully automatic chemistry analyzer, full blood count by automated cell analyzer, thyroid function tests by automated immunoassay by TOSOH machine, coomb's test and urine test for culture. A conjugated bilirubin of greater than 20% of total bilirubin was considered to be abnormal. Statistical analysis was made using graph pad prism. Data are shown as mean (S.D.).

RESULTS

According to history & other signs & symptoms of the new born, they were grouped into five classes. 24 cases belonged to physiological jaundice. Breast feeding/nonfeeding jaundice -84 cases, breast milk jaundice -5 cases, jaundice due

to prematurity -5 cases and pathological jaundice -6 cases(2 of neonatal sepsis, 2 of neonatal hypothyroidism, 1 of congenital biliary atresia & 1 of ABO incompatibility). The mean age of onset of jaundice in these patients was different in different groups (table 1). There is history of hyperbilirubinemia in siblings of 30% of cases. Klebsiella & pseudomonas was detected in blood culture of two cases conferring the diagnosis of

neonatal sepsis. TSH & T4 levels in neonatal hypothyroidism were highly abnormal. Reticulocyte count is highest in ABO incompatibility cases as hemolysis is maximum in these newborns. Conjugated bilirubin level & liver function tests were highly abnormal in case of congenital biliary atresia. Demographic profile & general data are summarized in table 1. Laboratory results are summarised in table 2.

Table-1: Demographic Profile & General Data of Newborns

Type of Jaundice	No of Cases	Sex	Age (Days)	Age of Onset (Days)	Birth Weight (Gram)	Delivery Vaginal/ Cesarean	Method of Feeding	Therapy for Icterus
Physiological	24	M-13 F-11	4.33 (1.09)	2.58 (0.72)	2870 (188.26)	V- 70.8% C- 29.2%	BF-91.66% Bot F – 8.33%	No therapy. Self resolution.
Breast Feeding	84	M-53 F- 31	6.6 (1.49)	3.19 (0.78)	2848 (150)	V-36.78% C-63.21%	BF-24% Bot F-33.3% MF -39%	Increase in frequency of BF & phototherapy.
Breast Milk	5	M- 4 F-1	22.4 (2.4)	16.6 (0.89)	2765 (155)	V – 60% C – 40%	BF– 80% MF – 20%	phototherapy
Prematurity	5	M- 3 F- 2	4.2 (0.84)	2.00 (0.71)	2092 (85.03)	V– 80% C – 20%	BF– 20% Bot F – 20% MF -60%	Phototherapy
Pathological								
Biliary Atresia	1	M-1	8	1	2468	C	BF	Surgical
Neonatal Sepsis	2	M -2	7.5 (0.71)	1.5 (0.71)	2395 (148.49)	V– 50% C – 50%	BF -50% MF– 50%	Antibiotic & phototherapy.
Neonatal Hypothyroidism	2	M-2	37.5 (10.61)	5.0 (1.41)	2900 (141)	V- 50% C– 50%	BF – 100%	Eltroxin & phototherapy
ABO Incompatibility	1	F-1	4	1	2568	V	BF	Photo therapy & Referral for exchange transfusion

Values are mean (S.D.), V- Vaginal Delivery, C-Cesarean Delivery, BF- Breast feeding, Bot F- Bottle feeding, MF- Mixed feeding

Table-2: Laboratory Investigations of Newborns

Type of Jaundice	Hb%	Reticulocyte Count	Highest Total Bilirubin mg/dl	Conjugated Bilirubin mg/dl	TSH uIU/ml	T4 ug/dl	ALT U/L	AST U/L	ALP U/L
Physiological	13.54 (1.25)	1.9 (0.42)	6.917 (0.67)	0.158 (0.072)	2.5 (0.41)	14.0 (1.38)	15.88 (4.07)	24 (6.7)	166.13 (53.28)
Breast Feeding	13.78 (1.25)	2.010 (0.30)	14.908 (1.96)	1.38 (0.22)	2.8 (0.34)	12.35 (0.88)	19.70 (4.41)	21.43 (7.16)	152.08 (50.41)
Breast Milk	11.40 (2.07)	1.48 (0.47)	15.20 (1.92)	1.88 (0.497)	2.72 (0.26)	11.6 (1.14)	21.6 (4.28)	25.6 (2.3)	92.6 (17.34)
Prematurity	10.00 (0.71)	1.14 (0.27)	7.6 (1.14)	0.66 (0.13)	1.32 (0.37)	12.2 (0.84)	14.8 (2.77)	19.6 (2.30)	58.2 (3.19)
Pathological Jaundice									
Neonatal Sepsis	8.5 (0.71)	5.00 (1.41)	17.00 (1.41)	0.35 (0.07)	0.9 (0.14)	8.5 (0.71)	41 (1.41)	49 (1.41)	67 (4.24)
Biliary Atresia	13	3	18	16	3.0	13	50	63	4000
ABO Incompatibility	14	9	23	2.4	3.0	13	15	29	160
Neonatal Hypothyroidism	12 ± 0.0	0.95(0.07)	14 (1.41)	2.05 (0.21)	47 (9.90)	1.57 (0.18)	16.5 (0.71)	28 (2.8)	120 (7.1)

Values are mean (S.D.), The neonate ,whose bilirubin was high because of ABO incompatibility , had blood group A-ve, and that of mother's was B+ve & this baby was the second issue of the lady

DISCUSSION

In current study, breast feeding/non feeding jaundice emerged as the most common aetiology of neonatal jaundice. Bilirubin level regressed after increasing the frequency & improving the method of breast feeding. Najati et al.,(2010)^[15] also concluded that majority of patients had an unconjugated hyperbilirubinemia probably due to breast non feeding. Other causes in their study were G6PD deficiency, hypothyroidism, UTI, septicaemia, Down syndrome & ABO incompatibility. Our study also has few cases of these clinical conditions as a cause of neonatal hyperbilirubinemia. Gartner LM (2001)^[16] stated in his study that insufficient caloric intake resulted from maternal and/or infant breast feeding difficulties may also increase unconjugated serum bilirubin concentration. Another study by Bertini (2001)^[17], stated that fasting plays an important role in pathogenesis of neonatal hyperbilirubinemia & forms bulk of the cases.

Breast milk jaundice was the diagnosis of exclusion as also stated by Prashant GD.^[18] All 5 newborns were exclusively breast fed. Icterus started developing after 1st week of life & peak occurred at 3rd week (12-15mg/dl) & thereafter it regressed by its own requiring no treatment. This type is not a clinical disorder but recognised to be a normally occurring extension of physiological jaundice of new born and breast feeding should not be interrupted. Schneider API^[19] also concluded in his study that breast milk jaundice is one of the common causes of neonatal jaundice.

Prematurity was another reason for neonatal hyperbilirubinemia. 5 cases were there. These newborns are prematurely delivered at the mean gestational age of 32 ± 2 weeks. 1 patient out of 5 showed neurological sequelae depicting the diagnosis of kernicterus.^[20] Onyearugha et al., (2011)^[21], concluded prematurity as the second leading cause of neonatal jaundice. According to another study, these neonates fed on human milk had higher peak concentration of plasma

bilirubin & more prolonged hyperbilirubinemia than those fed on an artificial infant formula.^[22]

Two cases of neonatal sepsis were reported. Blood culture showed presence of klebsiella & pseudomonas respectively. Extensive chemotherapy was instituted resulted in regression of jaundice. Onyearugha et al., (2011)^[21] also found sepsis as the second leading cause of jaundice in neonates in Nigeria.

Two cases of neonatal hypothyroidism was noted. Scott et al.,(2004)^[23] & Najati et al.,(2010)^[15] concluded in their study that hypothyroidism led to neonatal hyperbilirubinemia. ABO incompatibility & congenital biliary atresia also contributed as etiologies of hyperbilirubinemia in neonates.

Careful education about breast feeding & monitoring of mothers as well as assessment of newborns for the risk of developing hyperbilirubinemia can aid in preventing neonatal jaundice.

Treatment is based on total serum bilirubin concentration 6 hourly during phototherapy & exchange transfusion.

CONCLUSION

The present study concludes that breast feeding/nonfeeding is the most common cause of neonatal hyperbilirubinemia in areas near Shri Ram Murti Smarak Institute Of Medical Sciences, a tertiary care teaching hospital, followed by other causes like breast milk jaundice, jaundice due to prematurity, neonatal sepsis, neonatal hypothyroidism, Congenital biliary atresia & ABO incompatibility.

ACKNOWLEDGEMENT

Authors are thankful to guardians of newborns for their cooperation in history and examination of the neonate. Authors are also in debt of department of Paediatrics for their support in making diagnoses of neonatal jaundice & management plan of the newborn.

REFERENCES

- Slusher TM, Angyo IA, Bode TF. Transcutaneous bilirubin measurements and serum total bilirubin levels in indigenous African infants. *Pediatrics* 2004; 113: 1636-1641.
- Haque KM, Rahman M. An unusual case of ABO hemolytic disease of the new born. *Bangladesh Med Res Counc* 2000; 16: 105-108.
- Madan A, James RM, Stevenson DK. Neonatal Hyperbilirubinemia. In: Taeusch HW, Ballard RA, Gleason CA. *Avery's diseases of the new born*. 8th Ed. Elsevier Saunders 2004: 1226-1256.
- Laforgia N, Faienza MF, Rinaldi A. Neonatal hyperbilirubinemia and gilbert's syndrome. *J Perinat Med* 2002; 30: 166-169.
- Wang M, Hays T, Ambruso DR. Hemolytic diseases of the new born caused by a high titer antigroup B Ig G from a group A mother. *Pediatr Blood cancer* 2005; 45(6): 861-862.
- Ho NK. Neonatal jaundice in Asia. *Baillieres. Clin Haematol* 2002; 5: 131-142.
- Barbara JS, Kliegman RM. Jaundice and Hyperbilirubinemia in the Newborn. In: Kliegman RM, Behrman HB, Jenson HB. *Nelson Textbook of Pediatrics*, 17th Ed. Elsevier Saunders 2004: 592-598.
- Maisels MJ. Neonatal Hyperbilirubinemia. In: Klaus MH, Fanaroff A. *Care of the high risk neonate*. 5th Ed. Saunders company 2001: 324-341.
- Maria FB, Cecilia MD. Neonatal Jaundice and breast feeding. *Neo Reviews* 2007; 8: 282-288.
- Weissman LE, Merenstein GB, Digirol M. The effect of early meconium evacuation on early onset hyperbilirubinemia. *Am J Dis Child* 1988; 137: 666-8.
- Broderson R, Herman LS. Intestinal absorption of unconjugated bilirubin: a possible contributing factor in neonatal jaundice. *Lancet* 1963; I: 1242.
- Cynthia RH. Breast milk and breast feeding jaundice. Section 1 Diseases and Disorders. In: Lynne CG, Jeffrey K, Cynthia C. *Mosby's pediatric clinical advisor: instant diagnosis & treatment*. 2nd Ed. Elsevier 2002: 200.
- Kumal A, Ozkan H, Duman N. Breast milk jaundice correlates with high levels of EGF. *Pediatr Res* 2009; 66: 218-21.
- Mishra S, Agarwal R, Deorari AK. Jaundice in the new borns. *Indian J Pediatr* 2008; 75: 157-163.
- Najati N, Gharebaghi MM, Moratazavi F. Underlying etiologies of prolonged icterus in neonates. *Pakistan Journal of Biological Sciences* 2010; 13: 711-14.
- Gartner LM. Breastfeeding and Jaundice. *Perinatol* 2001; 1: S25-9.
- Bertini G, Dani C, Tronchin M. Is breast feeding really favoring early neonatal jaundice? *Paediatrics* 2001; 107: E41.
- Prashant GD. Breast milk jaundice work up. *E medicine. medscape. com/article/973629* 2010
- Schneider AP 2nd. Breast milk jaundice in the newborn. A real entity. *JAMA* 1986; 255: 3270-4.
- Camille M, Smith MD, Galen P, Barns BS, Claire A, Jacobson AD, et al. Auditory brainstem response response detects early bilirubin neurotoxicity at low indirect bilirubin values. *Journal of Perinatology* 2004; 24: 730-32.
- Onyearugha CN, Onyire BN, Ugboma HA. Neonatal Jaundice: Prevalence and associated factors as seen in Federal Medical Center Abakaliki, Southeast Nigeria. *Journal of Clinical Medicine and Research* 2010; 3: 40-45.
- Lucas A, Baker BA. Breast milk jaundice in premature infants. *Arch Dis Child* 1986; 61: 1063-67.
- Scott R, Aldangady N, Maalouf E. Neonatal hypopituitarism presenting with poor feeding, hypoglycaemia & prolonged unconjugated hyperbilirubinemia. *J Matern Fetal Neonatal Med* 2004; 16: 131-133.

Cite this article as: Kamal S, Sayedda K, Ahmed Q. Breast non feeding: Main cause of neonatal hyperbilirubinemia in areas adjoining Shri Ram Murti Smarak Institute of Medical Sciences, a tertiary care teaching hospital, Bareilly. *Natl J Physiol Pharm Pharmacol* 2012; 2:108-112.

Source of Support: Nil

Conflict of interest: None declared